

POSTOPERATIVE INSTRUCTIONS

ACHILLES TENDON REPAIR

DIET

- Begin with clear liquids and light foods (jellos, soups, etc.)
- Progress to your normal diet if you are not nauseated

WOUND CARE

- To avoid infection, keep surgical incisions clean and dry – you may shower by placing a large garbage bag over your splint starting the day after surgery – NO immersion of operative leg (i.e. bath)

MEDICATIONS

- Pain medication is injected into the wound and ankle joint during surgery – this will wear off within 8-12 hours
- Most patients will require some narcotic pain medication for a short period of time – this can be taken as per the directions on the bottle
- Common side effects of the pain medication are nausea, drowsiness, and constipation – to decrease the side effects, take medication with food – if constipation occurs, consider taking an over-the-counter laxative
- If you are having problems with nausea and vomiting, contact the office and ask for **Dr. El-Gazzar** at **newedgeorthopedics@gmail.com** or **(201) 985-8967**
- Do not drive a car or operate machinery while taking the narcotic medication
- Ibuprofen 200-400mg (i.e. Advil) may be taken in between the narcotic pain medication to help smooth out the post-operative ‘peaks and valleys’, reduce overall amount of pain medication required, and increase the time intervals between narcotic pain medication usage

ACTIVITY

- Elevate the operative leg to chest level whenever possible to decrease swelling
- Use crutches to assist with walking – use a heel-toe pattern when walking, but do NOT bear any weight on your operative leg – unless instructed otherwise by physician
- Do not engage in activities which increase ankle pain/swelling (prolonged periods of standing or walking) over the first 7-10 days following surgery
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks
- NO driving until instructed otherwise by physician
- May return to sedentary work ONLY or school 3-4 days after surgery, if pain is tolerable

ICE THERAPY

- Begin immediately after surgery
- Use icing machine continuously or ice packs (if machine not prescribed) every 2 hours for 20 minutes daily until your first post-operative visit – remember to keep leg elevated to level of chest while icing

EXERCISE

- No exercise or motion is to be done until instructed to do so by your physician after the first post-operative visit
- Formal physical therapy (PT) will begin about 10-14 days post-operatively with a prescription provided at your first post-operative visit

FOR EMERGENCY

- Contact **Dr. El-Gazzar** or his office **New Edge Orthopedics** at **newedgeorthopedics@gmail.com** or **(201) 985-8967** if any of the following are present:
 - Painful swelling or numbness
 - Unrelenting pain
 - Fever (over 101° - it is normal for the first day or two following



surgery) or chills

- Redness around incisions
- Color change in wrist or hand
- Continuous drainage or bleeding from incision (a small amount of drainage is expected)
- Difficulty breathing
- Excessive nausea/vomiting

